

# ***Observation of Structures in the $J/\Psi\phi$ from $B^+ \rightarrow J/\psi\phi K^+$ Decay at CMS***



***Kai Yi***

***University of Iowa***


***LBL RPM, Dec 20, 2012***

# *Outline*


- *Introduction to exotic mesons*
- *Motivation for  $J/\Psi\phi$  structures*
- *Status of  $J/\Psi\phi$  before CMS result*
- *CMS detector and trigger*
- *$J/\Psi\phi$  Analysis strategy and event selection*
- *Results*
- *Summary*

# Quark Model


- *The birth of quark model (M. Gell-Mann & G. Zweig):  
M. Gell-Mann, Phys. Lett. 8, 214 (1964)*
- *Heavy top decays before forming bound states  
light quarks exist as bound states*



– Baryons:  $(qqq)$



– Mesons:  $(q\bar{q})$   
quarkonia:  $(s\bar{s}), (c\bar{c}) (b\bar{b})$  (hidden)



- *J/Ψ establishes the quark model, Υ(1S) further confirms it*
- *Gell-Mann also suggested exotic states  $(qqq\bar{q}\bar{q}), (qqq\bar{q}\bar{q})$  at the birth of quark model, but evidence has never been solidly established*

*Revitalized* by recently discovered charmonium-like states  
despite almost a decade, still mysterious!


# From strange to bottom discovery

$\Omega^-$  discovery

$J/\psi$  ( $\bar{c}c$ ) discovery


$\Upsilon$  ( $\bar{b}b$ ) discovery

BNL




1964

IN THE BEGINNING, .....




1968

BNL




1974

SLAC



1974

FNAL



1977

*Heavy flavor quarkonium spectroscopy helped turn quarks into a reality!*  
*What we can learn from quarkonium-like spectroscopy?*



# Charmonium ( $c\bar{c}$ ) Potential Model (Cornell Model)

- *simple QCD-inspired phenomenological potential :*

$$V(r) = -\frac{\kappa}{r} + \frac{r}{a^2}, \quad \kappa = 0.61, m_c = 1.84 \text{ GeV}, a = 2.38 \text{ GeV}^{-1}$$

- **non-relativistic** (charm quark is “heavy” compared to binding energy)
- quark confinement (increases linearly with separation)
- extendable to include **spin-dependent** terms, **relativistic** corrections, etc.
- Lattice QCD provides calculation of the masses and widths

[Eichten et. al., PRD 17, 3090 \(1978\)](#)

[Godfrey & Isgur, PRD 32, 189 \(1985\)](#)

[Barnes et. al., PRD 72, 054026 \(2005\)](#)

# Charmonium States

## Notation:

$$2^{S+1}[L]_J$$

**L=S,P,D** (0,1,2)  
(No cand. with  
 $L \geq 3$ )

$$\mathbf{J} = \mathbf{L} + \mathbf{S}$$

$$\mathbf{S}(q\bar{q}) = 0 \text{ or } 1$$

$$\text{Parity: } \mathbf{P} = (-1)^{L+1}$$

Charge conjugation  
eigenvalues:  
 $\mathbf{C} = (-1)^{L+S}$


**N:** Radial  
Quantum  
Numbers

Quantum numbers				Name	Mass (MeV/c <sup>2</sup> )	width(MeV)
N	L	J <sup>PC</sup>	N <sup>2S+1</sup> L <sub>J</sub>			
1	0	0 <sup>++</sup>	1 <sup>1</sup> S <sub>0</sub>	$\eta_c(1S)$	2980.4±1.2	26.7±3
1	0	1 <sup>--</sup>	1 <sup>3</sup> S <sub>1</sub>	J/ψ	3096.916±0.011	93.2±0.02 ×10 <sup>-3</sup>
1	1	0 <sup>++</sup>	1 <sup>3</sup> P <sub>0</sub>	$\chi_{c0}(1P)$	3414.75±0.31	10.2±0.7
1	1	1 <sup>++</sup>	1 <sup>3</sup> P <sub>1</sub>	$\chi_{c1}(1P)$	3510.66±0.07	0.89±0.05
1	1	2 <sup>++</sup>	1 <sup>3</sup> P <sub>2</sub>	$\chi_{c2}(1P)$	3556.20±0.09	2.03±0.12
1	1	1 <sup>+-</sup>	1 <sup>1</sup> P <sub>1</sub>	$h_c(1P)$	3525.93±0.27	<1
1	2	1 <sup>--</sup>	1 <sup>3</sup> D <sub>1</sub>	$\psi(3770)$	3772.92±0.35	27.3±1.0
2	0	0 <sup>++</sup>	2 <sup>1</sup> S <sub>0</sub>	$\eta_c(2S)$	3637±4	14±7
2	0	1 <sup>--</sup>	2 <sup>3</sup> S <sub>1</sub>	$\psi(2S)$	3686.09±0.04	317±9 ×10 <sup>-3</sup>
2	1	2 <sup>++</sup>	2 <sup>3</sup> P <sub>2</sub>	$\chi_{c2}(2P)$	3929±5	29±10
3	0	1 <sup>--</sup>	3 <sup>3</sup> S <sub>1</sub>	$\psi(4040)$	4039±1	80±10
2	2	1 <sup>--</sup>	2 <sup>3</sup> D <sub>1</sub>	$\psi(4160)$	4153±3	103±8
4	0	1 <sup>--</sup>	4 <sup>3</sup> S <sub>1</sub>	$\psi(4415)$	4421±4	62±20

*These states work well with charmonium model, until the appearance of X(3872)*


# Hints before the discovery of $X(3872) \rightarrow J/\psi \pi^+ \pi^-$

*CDF internal, 1994*




*E705, PRD 50, 4258 (1994)*

*E705 saw  $\psi(3836)$  in 1994?*



*BaBar internal, 2003*



*CDF saw a hint in 1994, unpublished*  
*BaBar saw a hint in 2003, unpublished*

*Both CDF and Babar spotted hints of  $X(3872)$  before its discovery!*

*The mass from E705's  $\Psi(3836)$  is low*


From BaBar B-Factory Symposium (C. Hearty)  
<http://www-conf.slac.stanford.edu/b-factory-symposium/talks.asp>



# X(3872)--2003

PRL 91, 262001

$N^{2S+1}L_J$	$J^{PC}$	$u\bar{d}, u\bar{u}, d\bar{d}$ $I = 1$	$u\bar{u}, d\bar{d}, s\bar{s}$ $I = 0$	$c\bar{c}$ $I = 0$
$1^1S_0$	$0^{-+}$	$\pi$	$\eta, \eta'$	$\eta_c(1S)$
$1^3S_1$	$1^{--}$	$\rho$	$\omega, \phi$	$J/\psi(1S)$
$1^1P_1$	$1^{+-}$	$b_1(1235)$	$h_1(1170), h_1(1380)$	$h_c(1P)$
$1^3P_0$	$0^{++}$	$a_0(1450)^*$	$f_0(1370)^*, f_0(1710)^*$	$\chi_{c0}(1P)$
$1^3P_1$	$1^{++}$	$a_1(1260)$	$f_1(1285), f_1(1420)$	$\chi_{c1}(1P)$
$1^3P_2$	$2^{++}$	$a_2(1320)$	$f_2(1270), f_2'(1525)$	$\chi_{c2}(1P)$
$1^1D_2$	$2^{-+}$	$\pi_2(1670)$	$\eta_2(1645), \eta_2(1870)$	
$1^3D_1$	$1^{--}$	$\rho(1700)$	$\omega(1650)$	$\psi(3770)$
$1^3D_2$	$2^{--}$			??
$1^3D_3$	$3^{--}$	$\rho_3(1690)$	$\omega_3(1670), \phi_3(1850)$	
$1^3F_4$	$4^{++}$	$a_4(2040)$	$f_4(2050), f_4(2220)$	
$2^1S_0$	$0^{-+}$	$\pi(1300)$	$\eta(1295), \eta(1440)$	$\eta_c(2S)$
$2^3S_1$	$1^{--}$	$\rho(1450)$	$\omega(1420), \phi(1680)$	$\psi(2S)$
$2^3P_2$	$2^{++}$	$a_2(1700)$	$f_2(1950), f_2(2010)$	
$3^1S_0$	$0^{-+}$	$\pi(1800)$	$\eta(1760)$	



$X(3872) \rightarrow J/\psi \pi^+ \pi^-$   
 $M = 3871.8 \pm 0.7 \pm 0.4 \text{ MeV}$   
 $\Gamma < 3.5 \text{ MeV @ 90\% CL}$

*(Problematic) features*

mass  $\sim 70 \text{ MeV} > 1^3D_2$  charmonium,  $J^{PC} = 1^{++}$  or  $2^{-+}$   
 $M(\pi^+ \pi^-)$  peaks as a  $\rho$ ,  $C=+$ , isospin=1 (charmonium--0)  
 Decays to  $J/\psi \gamma$  &  $\Psi' \gamma$ , suppressed for  $2^{-+}$


Mass close to  $DD^*$ , molecule is speculated  
 No charged partners observed, tetra-quark?  
 similar rate as charmoniums at hadron colliders.  
 mixture of a  $DD^*$  molecule and the  $2^3P_1$  charmonium?

First particle challenging charmonium model, revitalized exotic meson study




# $Y(3940) \rightarrow J/\psi \omega$ , $Y(4260) \rightarrow J\psi \pi^+ \pi^-$ -- 2005

PRL 94, 182002



PRL 95, 142001




Above  $D\bar{D}$  &  $DD^*$  threshold,  
 Tiny branching fraction expected  
 New mass and width from BaBar:  
 $M \approx 3919.1^{+3.8}_{-3.4} \pm 2.0$ ,  $\Gamma \approx 31^{+10.8}_{-8} \pm 5 \text{ MeV}$   
[arXiv:1012.0074 \[hep-ex\]](https://arxiv.org/abs/1012.0074)  
 at the  $J/\psi \omega$  threshold ?

Well above  $D\bar{D}$  &  $DD^*$  threshold,  
 Tiny branching fraction expected  
 $J^{PC} = 1^-$ , plus  $Y(4350)$ ,  $Y(4660)$   
 too many  $1^-$  ?

# $Z(4430)^+ \rightarrow \psi(2S)\pi^+$ -- 2008

PRL 100, 142001



The first *charged charmonium-like* state, a smoking gun if confirmed


*Babar disagrees with Belle*

*Many more new states...*

*They do not fit into charmonium expectation  
Has been extended to bottomonium system*

*Beyond  $(q\bar{q})$  mesons: **exotic mesons?***

# Exotic Models-I



## Molecular

Loosely *bound state* of a pair of *mesons*. The dominant binding mechanism should be *pion exchange*. Being weakly bound the mesons tend to decay as if they were free



## Tetraquark

Bound state of *four quarks*, i.e.  $qq\bar{q}\bar{q}$  in which the quarks group into color triplet scalar or vector clusters

Strong decays proceed via rearrangement processes

***Distinctive features of multi-quark picture with respect to charmonium:***

- *prediction of many new states*
- *possible existence of states with non-zero charge, strangeness or both*




# Exotic Models-II



## Charmonium hybrids

States with excited gluonic degrees of freedom;  
exotic  $J^{PC}=0^{+-}, 1^{-+}, 2^{+-}$  ... not allowed for charmonium.  
Smoking gun for exotic states.

Lattice QCD for  $1^{-+}$ :  $m \sim 4.3 \pm 0.05 \text{ GeV}$  (C. Thomas)



Threshold, cusp, or coupled-channel effect giving a cross section enhancement which may not correspond to resonance production at all

## Hadro-charmonium

Light hadrons bounded by *van der Waal's force* to a charmonium core in the case where the light hadron is a highly excited resonance.

We know something is going on even though we do not know exactly what!

**New kind(s) of spectroscopy with complex binding forces?**

How about  **$J/\psi\phi$  system?** (threshold @4.116 GeV, VV, C=+)

(cc) with a mass above 4.116 GeV, expect **tiny** branching fraction to  $J/\psi\phi$ .

# Charmonium hybrid $\rightarrow J/\psi\phi$ ?

$J^{PC}$	Open charm	Hidden charm
$0^{+-}$	Quantum numbers forbid $D^{(*)}D^{(*)}$	$J/\psi\{f_{\{0,1,2\}}, (\pi\pi)_S\}$ $h_c\eta; \eta_c h_1$ $\chi_{c0}\omega$ $\chi_{c\{1,2\}}\{\omega, h_1, \gamma\}$
$0^{--}$	$D^*D$	$h_c(\pi\pi)_S$ $J/\psi\{f_{\{1,2\}}, \eta^{(')}\}$ $\chi_{c0}h_1; \eta_c\{\omega, \phi\}$ $\chi_{c\{1,2\}}\{\omega, h_1, \gamma\}$
<b><math>1^{-+}</math></b>	$D^*D, D^*D^*$	$\chi_{c\{0,1,2\}}(\pi\pi)_S$ $\eta_c\{f_{\{1,2\}}, \eta^{(')}\}$ $\chi_{c\{1,2\}}\eta$ $\{h_c, J/\psi\}\{\omega, h_1, \phi, \gamma\}$
$2^{+-}$	$D^*D, D^*D^*$	$\{h_c, J/\psi\}\{f_{\{0,1,2\}}, (\pi\pi)_S\}$ $\{h_c, J/\psi\}\eta^{(')}$ $\{\eta_c, \chi_{c\{0,1,2\}}\}\{\omega, h_1, \phi, \gamma\}$

← PRD 57, 5653 (1998)  
F. Close et al

Accessible at CMS

Considered to be the ground exotic state, mass prediction from 3.9 to 5.3 GeV  
Most recent Lattice QCD calculation:  $4.3 \pm 0.05$  GeV

# Multi-quark states $\rightarrow J/\psi\phi$ ?

$J^{PC}$	M(MeV)	Decay Channel
$0^{++}$	3834	-
$0^{++}$	3927	$J/\psi \omega$
$0^{-+}$	4277(+15)	$J/\psi \phi, J/\psi \omega, D_s^{*+} D_s^{*-}$
$0^{-+}$	4312(+30)	$J/\psi \phi, J/\psi \omega, D_s^{*+} D_s^{*-}$
$0^{--}$	4297(-5)	$\psi \eta(\eta'), D_s^+ D_s^-$
$1^{++}$	3890	$J/\psi \omega$
$1^{+-}$	3870	$J/\psi \eta$
$1^{+-}$	3905	$J/\psi \eta$
$1^{-+}$	4321(+15)	$J/\psi \omega, J/\psi \phi$
$1^{-+}$	4356 (+30)	$J/\psi \omega, J/\psi \phi$
$1^{--}$	4330	$\psi \eta(\eta'), D_s^{(*)+} D_s^{(*)-}; J/\psi f_0(980)$
$1^{--}$	4341(-5)	$\psi \eta(\eta'), D_s^{(*)+} D_s^{(*)-}; J/\psi f_0(980)$
$1^{--}$	4390(+40)	$\psi \eta(\eta'), D_s^{(*)+} D_s^{(*)-}; J/\psi f_0(980)$
$1^{--}$	4289(-41)	$\psi \eta(\eta'), D_s^{(*)+} D_s^{(*)-}; J/\psi f_0(980)$

← [arXiv:0902.2803](https://arxiv.org/abs/0902.2803)  
N. V. Drenska et al

$J/\psi\phi$  is well motivated!

How to search?


Inclusive? Challenge!

Through B decays!




# Search structures $\rightarrow J/\psi\phi$ through $B$ decays

- Experimentally attractive to search through clean  $B \rightarrow J/\psi\phi K$  channel
  - taking advantage of  $B$  lifetime and narrow  $B$  mass window
  - $B \rightarrow J/\psi\phi K$  is OZI suppressed, so low rate from phase space decays




vacuum polarization



gluon coupling


# The status before Original CDF Report

- The *status* through  $B \rightarrow J/\psi \phi K$  :



$J/\psi \rightarrow \mu\mu$  and  $ee$ , 10  $B^+$  and  $B^0$



*BaBar* 2003, PRL 91, 071801



$J/\psi \rightarrow \mu\mu$  and  $ee$

- statistically limited, no structures reported



# First Report by CDF w/ 2.7 fb<sup>-1</sup> (2009)



PRL 102:242002, 2009

Purity ~80% in B<sup>+</sup> region

Nice  $\phi$  shape



Near threshold peak, called Y(4140)

Significance: ~4 $\sigma$

Yield = 14 ± 5

$M = 4143.0 \pm 2.9$  (stat) ± 1.2 (syst) MeV

$\Gamma = 11.7^{+8.3}_{-5.0}$  (stat) ± 3.7 (syst) MeV

Not likely to be charonium:



High mass w/ narrow width

Dalitz plot

$\Delta M = m(\mu^+ \mu^- K^+ K^-) - m(\mu^+ \mu^-)$



# Update from CDF w/ 6.0 fb<sup>-1</sup> (2010)



[arXiv:1101.6058 \[hep-ex\]](https://arxiv.org/abs/1101.6058)

$Yield_1 = 19 \pm 6; >5\sigma$

$M_1 = 4143.4^{+2.9}_{-3.0} (stat) \pm 0.6 (syst) \text{ MeV}$

$\Gamma_1 = 15.3^{+10.4}_{-6.1} (stat) \pm 2.5 (syst) \text{ MeV}$

$Yield_2 = 22 \pm 8; 3.1\sigma$


$M_2 = 4277.4^{+8.4}_{-6.7} (stat) \pm 1.9 (syst) \text{ MeV}$

$\Gamma_2 = 32.3.7^{+21.9}_{-15.3} (stat) \pm 7.6 (syst) \text{ MeV}$


$\chi^2/\text{dof}$  between old and new  $\Delta m$  is 7.2/3,  $p$ -value=6.5% w/ four regions

$$\frac{B(B^+ \rightarrow Y(4140)K^+, Y(4140) \rightarrow J/\psi\phi)}{B(B^+ \rightarrow J/\psi\phi K^+)} = 0.149 \pm 0.039(\text{stat}) \pm 0.034(\text{syst})$$

# Belle: Confirm or Refute? (2009, 2010)



Y(4140): 7.5 +4.9/-4.4 events  
 Statistical significance: 1.9 $\sigma$   
 Signal could not be identified.



Note: CDF and Belle do not contradict each other.  
 In Belle, B meson at rest on  $\Upsilon(4S)$  rest frame, Kaon momentum from  $\phi$  decay is low, especially just above  $J/\psi\phi$  threshold  
 → lower reconstruction efficiency.

## Summary



- B factories suffer from low  $p_T$  track efficiency
- Belle *cannot confirm or deny* the existence of Y(4140)
- Tevatron edge over B factories:  
 Low  $p_T$  kaons are boosted from B momentum

*no verdict from Babar*

Kenkichi Miyabayashi  
 (Nara Women's Univ.)  
 2010 May QWG7

*QWG 2010*

# LHC<sub>b</sub>: Contests CDF Report (2011)



LHC<sub>b</sub> confirms neither structure(s)  
 2.4σ disagreement with CDF measurement  
 @90% CL:

$$\frac{\mathcal{B}(B^+ \rightarrow X(4140)K^+) \times \mathcal{B}(X(4140) \rightarrow J/\psi\phi)}{\mathcal{B}(B^+ \rightarrow J/\psi\phi K^+)} < 0.07.$$


$$\frac{\mathcal{B}(B^+ \rightarrow X(4274)K^+) \times \mathcal{B}(X(4274) \rightarrow J/\psi\phi)}{\mathcal{B}(B^+ \rightarrow J/\psi\phi K^+)} < 0.08$$

## LHCb Versus CDF: Two Punches In The Face!

By Tommaso Dorigo | July 27th 2011 05:48 AM | 10 comments | [Print](#) | [E-mail](#) | [Track](#)


The first is the tentative observation of a new hadron, called Y(4140), a bump observed in the invariant mass of pairs of

result. Note that, as reported in the figure, if the CDF signal were as estimated by CDF, LHCb would have been able to fit 39+−9+−6 events. The Y(4140) is on very shaky ground at the moment, and the new PDG will likely change its status in the particle zoo... This is punch number 1.





# *LHC<sub>b</sub>: Contests CDF Report (2011)*



*LHCb is specifically designed to select Bottom/charm/exotic-quarkonium particles and the products of their decays*

*Excellent lepton and hadron Identification*


*Excellent mass resolution*

*~2.5 MeV for X(3872) decays*

*BUT LHCb did not confirm the existence of Y(4140). A serious challenge!*


*A result from a 3<sup>rd</sup> experiment is important!*

*For instance, CMS?*





# The CMS Detector




*Relevant sub-detectors*

# CMS Detector Performance

## Excellent muon/silicon detectors for quarkonium:

- Muon system
  - High-purity muon identification
  - Good dimuon mass resolution ( $\Delta m / m \sim 0.6\%$  for  $J/\psi$ )
- Silicon Tracking detector
  - excellent track momentum resolution ( $\Delta p_T / p_T \sim 1\%$ )
  - excellent vertex reconstruction and impact parameter resolution




## LHC luminosity and CMS trigger:

- collect data at increasing instantaneous luminosity
  - about  $5\text{fb}^{-1}$  from 2011 data at  $\sqrt{s}=7\text{ TeV}$   
(used for this analysis)
- Triggers are essential ingredients
  - Special trigger for different analysis


For this analysis:

displaced dimuon vertex &  
minimum (di)muon transverse momentum




# Analysis strategy (CMS 2012)

- I) Reconstruct  $B^+$  as:



- II) Search for structure in  $J/\psi \phi$  mass spectrum inside  $B^+$  mass window



<https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsBPH11026>

CMS: BPH-11-026



# Event Selections (CMS 2012)

-- $|\eta|$  for all tracks  $\leq 2.4$

--*probability( $\chi^2$ ) for  $J/\psi$  vertex fit  $> 10\%$ , probability( $\chi^2$ ) for  $B^+$  vertex fit  $> 1\%$*

-- $p_T(\text{kaon track}) > 1 \text{ GeV}$

-- *$J/\psi$  vertex flight length significance  $\geq 3$*

&

*Dataset A:  $p_T(J/\psi) > 7 \text{ GeV}$*

*Dataset B:  $p_T(J/\psi) > 7 \text{ GeV}$  &  $p_T(\mu^+/\mu^-) > 4 \text{ GeV}$*

--mass window:

*$J/\psi$  ( $\pm 150 \text{ MeV}$ ) and  $\phi$  in  $[1.008, 1.035] \text{ GeV}$  (Breit-Wigner shape)*


*constraint  $\mu^+\mu^-$  to  $J/\psi$  PDG mass value*

*Requirements are not optimized to be unbiased, confirm trigger requirements*



# The B Signal (CMS 2012)

$B^+ \rightarrow J/\psi \phi K^+$  decay



*Signal PDF: Gaussian*

*Background PDF:  
2<sup>nd</sup>-order Chebyshev polynomial*

*Mass: consistent with PDG value*

*Width: consistent with simulation*



*Largest  $B^+ \rightarrow J/\psi \phi K^+$  sample collected in the world up to date*

*~20 times of CDF statistics ( $115 \pm 12$ ); ~7.2 of LHCb statistics ( $346 \pm 20$ )*

# $J/\Psi$ and $\phi$ Signal (CMS 2012)

$m(\mu^+\mu^-)$  before forming the  $B$  signal

The  $B^+$  sideband subtracted  $m(K^+K^-)$   
where  $m(J/\psi\phi K^+)$  is within  $\pm 3\sigma$  of  $m(B^+)$





- A clear and clean  $J/\Psi$  signal
- Nice  $\phi$  lineshape, consistent with PDG parameters
- $B(J/\Psi\phi K^+)$  dominates after  $\phi$  mass restriction

# $J/\psi\phi$ Invariant Mass Spectrum (CMS 2012)

- ▶ The mass difference  $\Delta m = m(\mu^+\mu^-K^+K^-) - m(\mu^+\mu^-)$  is used
- *Extracting the  $\Delta m$  spectrum*
  - *Divide the dataset into the 20 MeV  $\Delta m$  bins*
  - *Extract the number of B events for each  $\Delta m$  by fitting the  $J/\psi\phi K$  spectrum*
    - ◆ Means fixed to the PDG B mass
    - ◆ RMS fixed to the signal MC values
  - *Plot the B yield as a function of  $\Delta m$*

$p_T(J/\psi) > 7\text{GeV}$

$p_T(J/\psi) > 7\text{GeV} \ \& \ p_T(\mu) > 4\text{GeV}$





*correct the spectrum by efficiency before fitting*

*Relative efficiency over  $\Delta m$  : approx. flat*

# Background Shape Studies (CMS 2012)

The phase space Dalitz projection on  $m^2(J/\psi\phi)$   
*generated events (red)*  
 Vs  
*reconstructed events (black)*

Sideband subtracted KKK mass  
*Phase Space MC (red)*  
 Vs  
*data (black)*




*CMS detector does not produce peaks*  
*Also imply relative flat efficiency*

*Possible  $K^*(1770)$ ?*  
*Does it effect  $\Delta m$ ?*



# Background Shape Studies (CMS 2012)



*Event mixing to study the  $\Delta m$  shape*

*--  $J/\psi, \phi, K^+$  from different event*

*--  $\phi, K^+$  from the same event,  $J/\psi$  from different event. This is to get the impact on  $\Delta m$  from possible  $\phi K^+$  resonances*

*Require the  $J/\psi \phi K^+$  mass around  $B$  mass*

*Event-mixing  $\Delta m$  shapes are slightly distorted compared to three-body phase space*

*However, the possible effect is on high  $\Delta m$  region and the three-body phase space shape is more conservative at low  $\Delta m$  region where the two structures are observed.*

## *Preliminary $\phi K^+$ Resonances Studies (CMS 2012)*

- *Generated simple Dalitz plot for known  $K^*$  ( $\phi K^+$ ). No similar structures seen in  $m(J/\Psi\phi)$  from reflections of these known  $K^*$ .*
- *No evidence of structures or deviation from phase space background shape found in  $m(\phi K^+)$  mass distribution after removing the two structures in  $m(J/\Psi\phi)$  in the data*
- *Possible structure(s) in  $m(J/\Psi K^+)$ ? No evidence so far*
- *Possible interference ? Could affect lineshape parameters, do not expect much since no evident big signal(s)*
- *A full amplitude analysis is more suitable, but limited by statistics and high non-B combinatoric background w/ current dataset*
- *In progress...*

# Null- and Signal-hypothesis Fits

	Mass (MeV)	Signal Yield
First Peak	$1051.5 \pm 2.0$	$355 \pm 46$
Second Peak	$1220.0 \pm 3.0$	$445 \pm 83$


Background: 3-body phase space

Signal: S-wave relativistic Breit-Wigner functions convolved with a Gaussian resolution function


Significance:  $>5\sigma$  for 1st peak

evidence for 2nd peak


background + 1 signal



background only hypothesis



background + 2 signal




# Robust Checks

- Many checks to investigate the robustness of the two structures
  - ✱ Variations on selection cuts, different background and signal shapes, different  $\Delta m$  binning...

- ✱ Different Background-subtraction technique: *sPlot*



*sPlot* is a technique of background-subtraction by weighting each event based on observed signal to background ratio.





# Robust Checks

- ✳ All main requirements are varied step by step to investigate possible bias
- ✳ Each sideband-subtracted  $\Delta m$  distribution is compared to the default one
- ✳ No indication of bias was found  
one example with tighter cuts and purer  $B$  sample is shown below:




◆  $B$  purity  $\sim 60\%$  within  $\pm 1.5\sigma$  of  $m(B^+)$

◆ similar  $\Delta m$  spectrum

# Result

- ▶ The efficiency-corrected  $\Delta m = m(\mu^+\mu^-K^+K^-) - m(\mu^+\mu^-)$



	Mass (MeV)	Signal Yield
First Peak	$1051.5 \pm 2.0$	$355 \pm 46$
Second Peak	$1220.0 \pm 3.0$	$445 \pm 83$

$$m_1 = 4148.2 \pm 2.0(\text{stat.}) \pm 4.6(\text{syst.}) \text{ MeV}$$


$$m_2 = 4316.7 \pm 3.0(\text{stat.}) \pm 7.3(\text{syst.}) \text{ MeV}$$

- ▶ observed a  $J/\psi\phi$  structure at 4148 MeV with a significance greater than  $5\sigma$  confirms the existence of  $Y(4140)$  for the first time from another experiment  
 $CDF Y(4140): m = 4143.4^{+2.9}_{-3.0}(\text{stat}) \pm 0.6(\text{syst})$
- ▶ evidence for a second structure at  $\sim 4317$  MeV in the same mass spectrum

# What is it?

## Charmonium Spectrum

← 2<sup>nd</sup> peak  
← Y(4140)





- Well **above** charm pair threshold
- Expect **tiny** BF to  $J/\psi\phi$
- Does **not** fit into charmonium
- Close  $J/\psi\phi$  threshold like Y(3940)
- What is it? Molecule/hybrid/threshold CUSP?

# Possible Interpretation of the Structures?

<http://indico.ifj.edu.pl/MaKaC/materialDisplay.py?contribId=832&sessionId=19&materialId=slides&confId=11> (last page)

Sookyung Choi (Belle), EPS

- ❑ Possible  $J^{PC}$ : S-wave:  $0^{++}, 1^{++}, 2^{++}$   
P-wave:  $0^{-+}, 1^{-+}, 2^{-+}, 3^{-+}$
- ❑ Lattice QCD for  $1^{-+}$  ( $c\bar{c}g$ ):  $4.3 \pm 0.05$  GeV
- ❑  $M = 4316.7 \pm 3.0(\text{stat}) \pm 7.3(\text{syst})$  MeV  
(CMS 2<sup>nd</sup> structure)
- ❑ Can the 2<sup>nd</sup> structure be  $1^{-+}$  hybrid?  
expect to see it in  $J/\Psi\omega$  if so






- ❑ What is the 1<sup>st</sup> structure? The same kind or it can be a different kind compared to the 2<sup>nd</sup> one?  
Similar to  $Y(3940)$ , both close to  $VV$  threshold? Same kind?

A topic to be investigated & more in  $B^+ \rightarrow J/\Psi\phi K^+$



# Search Possible Charged Exotics in $J/\Psi K^+$ Spectrum

arXiv: 0708.3496 [hep-ph], Jonathan Rosner  
 Mechanism to produce  $Z^+(4430)$ :  
 $B \rightarrow K Z^+(4430), Z^+(4430) \rightarrow \Psi(2S)\pi^+$



Same mechanism, we can search for:

$$B^+ \rightarrow \phi Z^+(xxxx), Z^+(xxxx) \rightarrow J/\Psi K^+$$

## Summary







- *CMS observed two structures in the  $J/\psi\phi$  spectrum at 4148 MeV and 4317 MeV using  $5.2 \text{ fb}^{-1}$  of data at 7 TeV collision energy*

$$m_1 = 4148.2 \pm 2.0 \text{ (stat.)} \pm 4.6 \text{ (syst.) MeV } (>5\text{sigma})$$

$$m_2 = 4316.7 \pm 3.0 \text{ (stat.)} \pm 7.3 \text{ (syst.) MeV } (>3\text{sigma})$$




- *Confirm the existence of the  $Y(4140)$ , consistent with CDF result & find evidence for a second structure*
- *Preliminary investigation find no evidence of reflection from  $K^*$*
- *More to be expected with the large data sample (4X) from 2012*

*Stay tuned!*






*CDF/BELLE/LHCb are raw distributions w/o efficiency correction*

# Mini Summary of near $VV$ threshold behavior



PRD 77, 012001(2008)



$I(V,V)=0$ , observed near  $VV$  threshold enhancement, through (double) OZI suppressed process

$I(V)=1$ , no clear enhancement  
 Skip complicated  $\omega\omega, \rho\rho$

Observed near  $V(I=0)V(I=0)$  threshold enhancement. Strong decay. Above  $(qq'+q'q)$  threshold. What are they?



# Exotic $J^{PC}$

- For  $q\bar{q}$  meson system, let  $L$  to be the orbital angular momentum. The meson spin  $J$  is given by  $|L-S| < J < |L+S|$ , where  $S=0$  (antiparallel quark spin) or  $1$  (parallel quark spin)
- The parity  $P$  and charge parity  $C$  of the meson system can be expressed as:  
$$P = (-1)^{L+1}$$
$$C = (-1)^{L+S}$$
- In the configuration of  $P=(-1)^J$ ,  $S=1$ ,  $CP=+1$ ,  $\Rightarrow$   
**Exotic  $J^{PC}$**  (not allowed for  $q\bar{q}$  meson):  
 **$0^-$ ,  $0^+$ ,  $1^-+$ ,  $2^+$ , ...**  
But exotic mesons can have these  $JPC$  due to additional degree of freedom.
- Identify **exotic  $J^{PC}$**  is helpful to identify **exotic mesons**