Planck 2014 Constraints on the Cosmic Neutrino(-like) Background

Marius Millea
UC Davis
On Behalf of the Planck Collaboration

Special thanks: Lloyd Knox (advisor)

Berkeley RPM

Dec 18, 2014





Caveat: I can show you anything shown at the conferences. For a few plots I will have to use 2013 results, but no major impact to this talk otherwise.

Today

Dec 1 | Dec 8 | Dec 15 | Dec 22

Diligent work

Data comes in

Original planned release date

Planned delayed release date

Outline

- Planck
- Number of neutrino-like species $N_{\rm eff}$
  - BAO and H_0
  - Damping and acoustic oscillation phase shifts
  - Constraints on axions and axion-like particles (ALPs)
- Sum of neutrino masses Σm_ν
  - Gravitational lensing

What is Planck?



Planck in 2009

Planck in 2009

LFI: 30, 44, 70 GHz



HFI: 100, 143, 217, 353, 545, 853 GHZ





PLANCK

"T" map





Q map



U map







Planck-LFI 44 GHz

PLANCK



−700 μK 700

+Q

-Q

+U

-U



−20 μK 20



−20 μK 20

Planck-LFI 70 GHz



+Q

-Q

+U

-U





Planck-HFI 100 GHz



+Q

-Q

+U

-U





Note: Q/U no longer smoothed, so most of what you see is galaxy + noise

Planck-HFI 143 GHz



Planck-HFI 217 GHz



Planck-HFI 353 GHz



Beautifully Consistent Data

70 GHz



100 GHz



100 GHz – 70 GHz



Different detector technologies, different systematics


WMAP
PLANCK
-300
T (μK)
300

large-scale modes intermediate small-scale modes



Let’s decompose into band-limited maps and compare those

large-scale modes

intermediate

small-scale modes



=



+



+



Fluctuation power



<-- large-scale modes

small-scale modes -->

Filtered to keep: large scales

Planck 143 GHz



WMAP 94 GHz



Comparison with WMAP: what's new?

Filtered to keep: large scales intermediate scales

Planck 143 GHz

WMAP 94 GHz









Filtered to keep... large scales

intermediate scales

small scales

Planck 143 GHz

WMAP 94 GHz













Filtered to keep... large scales

intermediate scales

small scales

Planck 143 GHz

WMAP 94 GHz














Planck TT spectrum
2013
Need:
•Non-equilibrium recombination
•Neutrino interactions
•Helium (BBN consistent)
•Etc...
$\mathcal{D}_l = l(l+1)C_l/2\pi\ [\mu K^2]$
$\Delta\mathcal{D}_l\ [\mu K^2]$
Multipole l



preliminary

Previous TE and EE measurements (Crites et al. 2014)



Perturbations in the tightly coupled photon-baryon plasma are just propagating “sound” waves governed by:

$$\delta'' + \frac{1}{c_s^2}\ddot{\delta} = 0$$

$$\ddot{\tilde{\delta}} + k^2 c_s^2 \tilde{\delta} = 0$$

(Caveat: ignores gravitational potentials and electron mass)

Perturbations in the tightly coupled photon-baryon plasma are just propagating “sound” waves governed by:

$$\delta'' + \frac{1}{c_s^2}\ddot{\delta} = 0$$

$$\ddot{\tilde{\delta}} + k^2 c_s^2 \tilde{\delta} = 0$$

(Caveat: ignores gravitational potentials and electron mass)

Consider Fourier modes projected onto the last scattering surface:



Perturbations in the tightly coupled photon-baryon plasma are just propagating “sound” waves governed by:

$$\delta'' + \frac{1}{c_s^2}\ddot{\delta} = 0$$

$$\ddot{\tilde{\delta}} + k^2 c_s^2 \tilde{\delta} = 0$$

(Caveat: ignores gravitational potentials and electron mass)

Consider Fourier modes projected onto the last scattering surface:




position


e
Electrons see dipole
e
position


e
Electrons see dipole
e
e
position
Electrons see quadrupole
e


e
Electrons see dipole
e
e
e
position
Electrons see quadrupole
e
e



“E mode” polarization pattern



“E mode” polarization pattern



“E mode” polarization pattern



“B mode” polarization pattern (can't be sourced by density/velocity perturbations)



“E mode” polarization pattern



“B mode” polarization pattern (can't be sourced by density/velocity perturbations)



Polarization maps are just linear combinations of the above patterns, e.g.:



“E mode” polarization pattern



“B mode” polarization pattern (can't be sourced by density/velocity perturbations)



Polarization maps are just linear combinations of the above patterns, e.g.:





- E modes and T modes out of phase
- With E modes, we are learning about velocities

Previous TE and EE measurements (Crites et al. 2014)



TE


140
70
0
-70
-140
$\mathcal{D}_\ell^{TE}$ $[\mu K^2]$
10
0
-10
$\Delta\mathcal{D}_\ell^{TE}$ $[\mu K^2]$
30
500
1000
1500
2000
ℓ

EE


×1e-05
100
80
60
40
20
0
$\mathcal{C}_\ell^{EE}$ $[\mu K^2]$
4
0
-4
$\Delta\mathcal{C}_\ell^{EE}$ $[\mu K^2]$
30
500
1000
1500
2000
ℓ





However, there are low-level systematics (mostly T->P leakage) in the polarization data whose solution was not fully tested in time for 2014.

The TT results are robust. The TE,EE results are more tentative and we don't quote them in our “baseline” results.

Summary of 2014 Results

- ΛCDM still consistent with the data, no clear preference for extensions
- Mean value for τ (optical depth to reionization) shifts lower by 1σ compared to WMAP
- Power spectrum preference for low H_0 & high σ_8 persists
- Some dark matter interpretations of the AMS/Fermi/Pamela excess are tentatively ruled out
- 40σ detection of CMB lensing, 715 new SZ clusters, etc...
- Cosmic Neutrino-(like) Background

This talk

Outline

- Planck
- Number of neutrino-like species $N_{\rm eff}$
  - BAO and H_0
  - Damping and acoustic oscillation phase shifts
  - Constraints on axions and axion-like particles (ALPs)
- Sum of neutrino masses Σm_ν
  - Gravitational lensing

All Aspects of Cosmology are Touched by the Planck Results

Observation-related Examples:

- BAO-determined distance-redshift relation
- SDSS matter power spectrum
- Deep Lens Survey cosmic shear power spectrum
- Other CMB measurements (e.g. WMAP, SPT, and ACT)

Consistent*

- Cepheids + SNe for determining H_0
- CFHTLS cosmic shear power spectrum
- σ_8 inferred from cluster counts

Some tension*

*Assuming the ΛCDM model

All Aspects of Cosmology are Touched by the Planck Results

Observation-related Examples:

- BAO-determined distance-redshift relation
- SDSS matter power spectrum
- Deep Lens Survey cosmic shear power spectrum
- Other CMB measurements (e.g. WMAP, SPT, and ACT)

Consistent*

- Cepheids + SNe for determining H_0
- CFHTLS cosmic shear power spectrum
- σ_8 inferred from cluster counts

Some tension*

*Assuming the ΛCDM model


t=0
LS surface
r_s
D_A
θ_s



Planck:

$$\theta_s(z = 1090) = (0.59672 \pm 0.00035)\mathrm{deg}$$

SDSS-BOSS:

$$\theta_{\mathrm{BAO}}(z = 0.57) = (4.1 \pm 0.07)\mathrm{deg}$$

$$\theta_{\mathrm{BAO}} \equiv r_s(z_{\mathrm{drag}})/(D_A^2/H)^{1/3}$$

BOSS BAO, Riess et al. (2011) H_0 and Planck

- Good agreement with BOSS BAO
- 2.4σ tension with Riess et al. H_0



BOSS BAO, Riess et al. (2011) H_0 and Planck

- Update to 2014



BOSS BAO, Riess et al. (2011) H_0 and Planck

- More conservative Hubble prior?



BOSS BAO, Riess et al. (2011) H_0 and Planck



BOSS BAO, Riess et al. (2011) H_0 and Planck

$$\rho_{\rm rad} = \rho_\gamma + \rho_\nu = \left[1 + \frac{7}{8}\left(\frac{4}{11}\right)^{4/3} N_{\rm eff}\right]\rho_\gamma$$

LCDM:
$N_{\rm eff} = 3.046$

Riess et al. 2011 (68%)

Anderson et al. 2014 (68%)



Outline

- Planck
- Number of neutrino-like species $N_{\rm eff}$
  - BAO and H_0
  - Damping and acoustic oscillation phase shifts
  - Constraints on axions and axion-like particles (ALPs)
- Sum of neutrino masses Σm_ν
  - Gravitational lensing


t=0
LS surface
r_d
r_s
D_A
θ_d

How does measuring these two scales lead to an N_{eff} constraint?

- Physics is remarkably simple, laid out in Hou et al. (2013), Bashinsky & Seljak (2004), Hu & White (1997)

How does measuring these two scales lead to an N_{eff} constraint?

- Physics is remarkably simple, laid out in Hou et al. (2013), Bashinsky & Seljak (2004), Hu & White (1997)

$$r_s = \int_0^{t_*} c_s \frac{dt}{a} = \int_0^{a_*} \frac{c_s da}{a^2 H(a)}$$

How does measuring these two scales lead to an N_{eff} constraint?

- Physics is remarkably simple, laid out in Hou et al. (2013), Bashinsky & Seljak (2004), Hu & White (1997)

$$r_s = \int_0^{t_*} c_s \frac{dt}{a} = \int_0^{a_*} \frac{c_s da}{a^2 H(a)}$$

$$r_d^2 = \pi^2 \int_0^{a_*} \frac{da}{a^3 \sigma_T n_e H(a)}$$

How does measuring these two scales lead to an N_{eff} constraint?

- Physics is remarkably simple, laid out in Hou et al. (2013), Bashinsky & Seljak (2004), Hu & White (1997)

$$r_s = \int_0^{t_*} c_s \frac{dt}{a} = \int_0^{a_*} \frac{c_s da}{a^2 H(a)}$$

$$r_d^2 = \pi^2 \int_0^{a_*} \frac{da}{a^3 \sigma_T n_e H(a)}$$

$$\theta_s = \frac{r_s}{D_A}$$

How does measuring these two scales lead to an N_{eff} constraint?

- Physics is remarkably simple, laid out in Hou et al. (2013), Bashinsky & Seljak (2004), Hu & White (1997)

$$r_s = \int_0^{t_*} c_s \frac{dt}{a} = \int_0^{a_*} \frac{c_s da}{a^2 H(a)}$$

$$r_d^2 = \pi^2 \int_0^{a_*} \frac{da}{a^3 \sigma_T n_e H(a)}$$

$$\theta_s = \frac{r_s}{D_A}$$

$$\theta_d = \frac{r_d}{D_A}$$

How does measuring these two scales lead to an N_{eff} constraint?

- Physics is remarkably simple, laid out in Hou et al. (2013), Bashinsky & Seljak (2004), Hu & White (1997)

$$r_s = \int_0^{t_*} c_s \frac{dt}{a} = \int_0^{a_*} \frac{c_s da}{a^2 H(a)}$$

$$r_d^2 = \pi^2 \int_0^{a_*} \frac{da}{a^3 \sigma_T n_e H(a)}$$

$$\theta_s = \frac{r_s}{D_A}$$

$$\theta_d = \frac{r_d}{D_A}$$

$$\frac{\theta_s}{\theta_d} = \frac{r_s}{r_d} \sim \frac{1}{\sqrt{H(a)}} \sim \frac{1}{N_{\rm eff}^{1/4}}$$

How does measuring these two scales lead to an N_{eff} constraint?

- Physics is remarkably simple, laid out in Hou et al. (2013), Bashinsky & Seljak (2004), Hu & White (1997)

$$r_s = \int_0^{t_*} c_s \frac{dt}{a} = \int_0^{a_*} \frac{c_s da}{a^2 H(a)}$$

$$r_d^2 = \pi^2 \int_0^{a_*} \frac{da}{a^3 \sigma_T n_e H(a)}$$

$$\theta_s = \frac{r_s}{D_A} \qquad \theta_d = \frac{r_d}{D_A}$$

$$\frac{\theta_s}{\theta_d} = \frac{r_s}{r_d} \sim \frac{1}{\sqrt{H(a)}} \sim \frac{1}{N_{\rm eff}^{1/4}}$$

$$H(a)^2 = \frac{8\pi G}{3} \left[\rho_r(a) + \rho_m(a) + \rho_\Lambda(a)\right]$$

Neff affects the ratio of sound horizon to diffusion scale



$$\frac{\theta_s}{\theta_d} \sim \frac{1}{N_{\rm eff}^{1/4}}$$

To scale

Neff affects the ratio of sound horizon to diffusion scale



$$\frac{\theta_s}{\theta_d} \sim \frac{1}{N_{\rm eff}^{1/4}}$$

To scale

Light Degrees of Freedom

$N_{\rm eff} = 3.13 \pm 0.32$ (*Planck* TT+lowP)

$N_{\rm eff} = 3.15 \pm 0.23$ (*Planck* TT+lowP+ BAO)

with high-l polarisation

$N_{\rm eff} = 2.98 \pm 0.20$ (*Planck* TT,TE,EE+lowP)

$N_{\rm eff} = 3.04 \pm 0.18$ (*Planck* TT,TE,EE+lowP+BAO)

but …



Slide: Zhen Hou

Data points: Planck 2013 data



Slide: Zhen Hou

Data points: Planck 2013 data

$$r_d^2 = \pi^2 \int_0^{a_*} \frac{da}{a^3 \sigma_T n_e H(a)}$$

$$\text{fixing } \Omega_b h^2, a_{\rm eq}, \theta_s, \theta_d$$

adjust Y_p to keep this fixed



CMB & BBN in great agreement



D/H measurements



Outline

- Planck
- Number of neutrino-like species $N_{\rm eff}$
  - BAO and H_0
  - Damping and acoustic oscillation phase shifts
  - Constraints on axions and axion-like particles (ALPs)
- Sum of neutrino masses Σm_ν
  - Gravitational lensing

Constraints on axions and ALPs

(Millea, Knox, and Fields 2014, in prep)

- The CMB and BBN can jointly probe scenarios where particles decay in between the two epochs.
- We consider a generic particle, called an axion-like particle (ALP) which can decay to two photons.



Axions and axion-like particles (ALPs)

- ALPs are a general class of particles with two free parameters: m_ϕ and $\tau_{\phi\gamma}$
  - Typical in theories with SSBs, e.g. string axiverse, etc..

$$\mathcal{L} = \frac{1}{2} m_\phi^2 \phi^2 + \frac{g_{\phi\gamma}}{4} \phi F \tilde{F}$$

$$\tau_{\phi\gamma} = \frac{64\pi}{m_\phi^3 g_{\phi\gamma}^2}$$

- Axions are a type of ALPs where $g_{\phi\gamma}$ and m_ϕ are related
  - Well motivated theoretically to solve the strong CP problem.
- Relevant axions and ALPs here are heavy compared to "invisible" axion models
  - $m_\phi \sim 1\ \mathrm{MeV}$
  - $\tau_{\phi\gamma} \sim 1\ \mathrm{sec}$



Reaction is fast when $\Gamma > H$



Two-photon interaction



$\Gamma \sim g^2_{\phi\gamma}$

Reaction is fast when $\Gamma > H$

Primakoff Process



rate [1/time]

H(a)

$\Gamma \sim n_q \alpha g_{\phi\gamma}^2$

Two-photon interaction



scale factor, a

$\Gamma \sim g_{\phi\gamma}^2$

Reaction is fast when $\Gamma > H$

Primakoff fast

Two-photon fast

Decoupled



Out-of-equilibrium decay:



Two-photon fast

Primakoff fast

In-equilibrium decay:




N_{eff}^{CMB}
$\log_{10}(\tau_{\phi\gamma}[\mathrm{sec}])$
$\log_{10}(m_\phi\,[\mathrm{eV}])$
<2.44
=2.44
=3.046
3.046
2.837
2.628
2.419
2.209
2.000
A
$m_\phi = 10^6$ eV
$\tau_{\phi\gamma} = 10^5$ sec
B
$m_\phi = 10^7$ eV
$\tau_{\phi\gamma} = 10^{-2}$ sec
$\rho(a)a^4$
a

Previous cosmological constraints from Cadamuro & Redondo (2012) + collider + beam dump

Two axion models



“MeV-ALP window”

Updates:
- Planck 2013
- Cooke et al. 2014 D/H
- Aver et al. 2013 Yp
- KEKB collider
- More accurate spectral distorition calculation

Two axion models



"MeV-ALP window"
No more...

Unless...



Diluted down to ~3 by CMB epoch

Can have 4 or 5 neutrino-like species initially

Slide: Zhen Hou

Data points: Planck 2013 data

$\text{fixing}\ \Omega_b h^2, a_{\text{eq}}, \theta_s, \theta_d$



Slide: Zhen Hou

Data points: Planck 2013 data

$\text{fixing}\,\Omega_b h^2, a_{\rm eq}, \theta_s, \theta_d$



Slide: Zhen Hou

Data points: Planck data

$\text{fixing}\,\Omega_b h^2, a_{\text{eq}}, \theta_s, \theta_d$

Phase shift appears to be contributing to Planck constraint



(Follin, Knox, Millea, and Pan 2014, in prep)

Gravitational potentials

$$\ddot{\tilde{\delta}} + k^2 c_s^2 \tilde{\delta} = F(\Phi, \Psi)$$

Neutrinos → potentials → photons

- Neutrino perturbations propagate at a different speed, which cause a phase shift in the acoustic oscillations.

$$C_\ell \to C_{\ell+\delta\ell}$$

$$\delta\ell = A(N_{\rm eff}, N_{\rm eff}^{\rm fid}) f_\ell$$

For the first time, we directly measure this phase shift:



(A similar test of neutrino perturbations as constraints on c_{vis} and c_{eff})

Outline

- Planck
- Number of neutrino-like species $N_{\rm eff}$
  - BAO and H_0
  - Damping and acoustic oscillation phase shifts
  - Constraints on axions and axion-like particles (ALPs)
- Sum of neutrino masses Σm_ν
  - Gravitational lensing



$$T(\hat{n}) \rightarrow T(\hat{n} + \nabla\phi(\hat{n}))$$

$$\phi(\hat{n}) = -2\int_0^{\chi_*} d\chi \frac{f_K(\chi_* - \chi)}{f_K(\chi_*) f_K(\chi)} \Psi(\chi\hat{n}; \eta_0 - \chi)$$

Two ways to analyze lensing with Planck

- Beginning with the Planck 2013 results, we inferred neutrino mass dominantly through their effect on lensing



Lensing potential reconstruction



The smoothness of the temperature power spectrum



Tension persists in 2014 between the amount of lensing measured the two.

Minimum from neutrino oscillations



This manifests as a loosening of neutrino mass constraints upon addition of lensing reconstruction data.

Conclusion

LCDM survives Planck 2014. Some hints of extensions in combination with other probes.



Neutrino number constraints from damping and phase shifts.



MeV axion ruled out, or maybe plays a role in a more speculative scenario?



Neutrino mass constraints from lensing.



Extra slides


Planck TT+lowP
+lensing
+lensing+BAO
ΛCDM
+$N_{\rm eff}$
+$N_{\rm eff}$+Σm_ν
+$N_{\rm eff}$
+$N_{\rm eff}$+Σm_ν
+Σm_ν
+$N_{\rm eff}$+$m^{\rm eff}_{\nu,\,\rm sterile}$
+Σm_ν
+$N_{\rm eff}$+$m^{\rm eff}_{\nu,\,\rm sterile}$
σ_8
0.9
0.8
0.7
0.24
0.32
0.40
0.48
$\Omega_{\rm m}$
56
64
72
H_0

Forecast from SNOWMASS





Slide: Duncan Hanson

Gravitational Lensing



Slide: Duncan Hanson

Gravitational Lensing



Slide: Duncan Hanson



Neff and mnu


Planck+WP+highL
Planck+WP+highL+BAO
4.8
4.0
3.2
2.4
$N_{\rm eff}$
0.0
0.2
0.4
0.6
0.8
1.0
Σm_ν [eV]


0.5
1.0
2.0
5.0
10.0
4.5
4.0
3.5
$N_{\rm eff}$
0.0
0.6
1.2
1.8
2.4
$m^{\rm eff}_{\nu,\,\rm sterile}$ [eV]
0.136
0.128
0.120
0.112
0.104
0.096
0.088
$\Omega_c h^2$

CMB Polarization and Lensing Reconstruction



SPT-3G: A proposed 2500 sq. deg. Survey with a 3rd-generation polarization-sensitive focal plane.

Enabling a deflection angle power spectrum measurement as forecasted here and

$\sigma(\Sigma m_\nu) = 0.06$ eV

A curiosity: Low-L tension with LCDM



Planck and WMAP in the 2<L<50 region





WMAP

WMAP+(no L<30 TT)

BOSS BAO, Riess et al. (2011) H_0 and Planck LCDM

- Planck is in excellent agreement with BAO measurement, discrepant with Riess et al. H_0




Planck
WMAP9
Cepheids+SNela
Riess et al. (2011)
Carnegie HP
Freedman et al. (2012)
HST Key Project
UGC 3789
RXJ1131−1231
SZ clusters
60
65
70
75
80
85
Expansion rate today
H_0 [km s^{-1}Mpc^{-1}]

The deflection angle power spectrum




Planck+WP+highL
Planck+lensing+WP+highL
Planck+WP+highL (A_L)
Planck−lowL+highL+τprior
Planck−lowL+lensing+highL+τprior
Planck−lowL+τprior
P/P_{max}
Σm_ν [eV]
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.4
0.8
1.2
1.6
2.0

Consistency Tests Within Same Frequency





Consistency Tests Between Different Frequencies

- In units of μK, the CMB is the same at all frequencies
- This is a critical tests of galactic foreground cleaning, extra-galactic foreground modeling, and transfer functions



Effect of modeling choices and data selection



217 failed null test detectors removed

The three things which most significantly affect H0 or Ωm



β/β^{dipole}

Component along dipole direction

Components along two directions perpendicular to the dipole direction

We derive, in multiple ways, a $\beta=v/c$ that is consistent in magnitude and direction with what's required to explain the dipole.

No Primordial Non-Gaussianity, just as expected from “slow-roll” inflation

f_{NL}^{local} is a phenomenological measure of non-Gaussianity



Non-zero!

But some signal expected due to a 2nd-order effect of late-time evolution (not primordial)

After subtraction of late-time effect:

$f_{NL}^{local} = 2.7 +/- 5.8$

> 5σ detection of scale dependence of primordial fluctuations ==> time dependence during inflation



Best-fit scale-invariant (n_s =1) model


WMAP9 LCDM
WMAP9
ACT
SPT
Planck
$\ell(\ell+1)C_\ell/2\pi\ [\mu\mathrm{K}^2]$
10^3
10^2
0
500
1000
1500
2000
2500
ℓ


$\ell(\ell+1)C_\ell/2\pi\ [\mu\mathrm{K}^2]$
10^3
0
100
200
300
400
500
ℓ

Here ACT/SPT/Planck are all sample variance limited but Planck has much larger sky coverage





Finally, at around l=2000, ACT/SPT become a tighter constraint because their beam is smaller





CMB Constraints on σ_8, Ω_m



Planck measurements favor a shift in σ_8 and Ω_m Driven by:

- 1st/3rd acoustic peak power ratio
- Gravitational lensing in the CMB power spectrum (Ω_m goes down by ~1σ when A_{Lens} is free)

	WMAP7	WMAP7+SPT	Planck-CMB
σ_8	0.819 +/- 0.031	**0.795 +/- 0.022**	**0.829 +/- 0.012**
Ω_m	0.276 +/- 0.029	**0.250 +/- 0.020**	**0.315 +/- 0.016**

(WMAP7) Komatsu +2011
(SPT) Story+2012
Planck XX 2013
Planck XVI 2013

Planck Cosmology has ***profound*** mismatch with Cluster Abundance





$$\text{Cluster counts} \sim (\sigma_8)^{10}$$

Vikhlinin et al. 2009 (CCCP, X-rays)

Planck Cosmology has ***profound*** mismatch with Cluster Abundance



Cluster counts ~ $(\sigma_8)^{10}$

Vikhlinin et al. 2009 (CCCP, X-rays)

Tension exists for ***SZ, X-ray, Optical*** cluster surveys and other probes of structure



- ***SZ, X-ray, and optical cluster surveys all favor lower*** σ_8, Ω_m (Reichardt+13, Vikhlinin +09, Rozo+10, etc.)

- **Other probes of structure are consistent with clusters**:
  - Weak lensing surveys (e.g, CFHTLS, Kilbinger+13)
  - Redshift space distortions (Macaulay+13)
  - Planck CMB lensing power spectrum (PlanckXVII)

- *A neutrino mass of* $\Sigma m_\nu \sim 0.3$ *eV would relieve this tension.*

However, I think its still reasonable to question evidence for high σ_8, Ω_m ***from Planck CMB.***

Current and Future SZ Surveys

2012: SPTpol
~1600 detectors



2013: ACTpol
~3000 detectors



2016: SPT-3G
~15,000 detectors



2018: CCAT
~20,000 detectors



2020+: PRISM
CMB-Lens
~100,000 detectors



Current and Future SZ Surveys

2012: SPTpol
~1600 detectors

2013: ACTpol

	Start Date	Area (deg^2)	Depth (uK-arcmin)	$N_{clusters}$
SPTpol	2012	500	6	**~1000**
ACTpol	2013	4000/150	20/4	**~1000**
SPT-3G	2016	2500	2	**~10,000**
CCAT	2018	~20,000	15	**~5000**
CMB-Lens	2020+	~20,000	~1	**~150,000**
PRISM	2020+	All-sky	~1	**~1 million**

SPT-3G:
Projected B-mode Power Spectrum



- Neutrino Constraints:
  - ***$\delta(N_{eff}) = 0.06$***
  - ***$\delta(\Sigma m\nu) = 0.06$ eV***
- "De-lens" the CMB at large-angular scales and improve "***r***" Inflation constraint
  - ***$\delta(r) = 0.01$***

Credit: T. Crawford

PIXIE


Calibrator (580 mm)
Sky
A Beam
B Beam
V Polarization
H Polarization
A
B
Folding Flats
Primary Mirror
550 mm f/1.7
Secondary Mirrors
50 mm f/7.5
Polarizer A
Transfer 1
125 mm f/2.4
Split Polarizations
Polarizer B
Transfer 2
125 mm f/4.4
Mix Beams
Dihedral Mirror
Transfer 3
125 mm f/4.4
FTS
Mirror Motion
Inject Phase Delay
Polarizer C
Transfer 4
125 mm f/4.4
Re-Combine Beams
Polarizer D
Transfer 5
125 mm f/4.4
Split Polarizations
L
R
Transfer 6
125 mm f/2.0
Measure Fringes
Detectors


Wavelength
3 cm
3 mm
300 μm
50 μm
Intensity (W / m² / sr / Hz)
10⁻²⁰
10⁻²²
10⁻²⁴
10⁻²⁶
Dust
CMB
Unpolarized
CMB
E-Mode
Synch
CMB
B-Mode
PIXIE Sensitivity
10
100
1000
Frequency (GHz)





Spectral Distortions

LCDM makes a very precise prediction




WMAP9 LCDM
Neff=1
nrun=-0.05
WMAP9
$\ell(\ell+1)C_\ell/2\pi\ [\mu\mathrm{K}^2\ ]$
10^3
10^2
0
500
1000
1500
2000
2500
ℓ


WMAP9 LCDM
Neff=1
nrun=-0.05
WMAP9
ACT
SPT
$\ell(\ell+1)C_\ell/2\pi\ [\mu\mathrm{K}^2\ ]$
10^3
10^2
0
500
1000
1500
2000
2500
ℓ


WMAP9 LCDM
WMAP9
ACT
SPT
$\ell(\ell+1)C_\ell/2\pi\ [\mu\mathrm{K}^2\ ]$
10^3
10^2
0
500
1000
1500
2000
2500
ℓ


WMAP9 LCDM
WMAP9
ACT
SPT
Planck
$\ell(\ell+1)C_\ell/2\pi\ [\mu\mathrm{K}^2]$
10^3
10^2
0
500
1000
1500
2000
2500
ℓ



ΛCDM is a good fit!



ΛCDM is a good fit!

Possible contamination affecting damping: unresolved foregrounds

To scale

Resolved foregrounds:



Possible contamination affecting damping: unresolved foregrounds

To scale

Resolved foregrounds:



Unresolved foregrounds:
(Noise-free simulation)



Adding “highL” to help constrain foregrounds

SPT highL





correlation coefficient ~ .5

potential for 25% tighter sigma

Consistency with other CMB experiments

- Planck and WMAP are consistent (except an overall “calibration”)
- Planck and SPT are consistent
- There *are* seemingly large differences between parameters from the three, but there's no evidence of any systematics.
  - Therefore one should combine them all, in which case Planck tends to dominate the result.

Planck-SPT consistency



SPT team cross-corelated Planck maps with SPT maps on the SPT patch of sky



Planck-WMAP consistency



There are differences between Planck and WMAP that look something like a 2.5% rescaling

Planck-WMAP consistency



There are differences between Planck and WMAP that look something like a 2.5% rescaling

The 2.5% difference is absorbed almost entirely by the amplitude



Parameters from L<800

What at L>800 is causing Planck LCDM parameter shifts?



- Its lensing
- This is going to be very important for the Σm_ν constraint

Outline

- Planck
- ΛCDM, the standard model of cosmology, passes a precision test
- Consistency with other cosmological probes
  - BAO and H0
  - WMAP and SPT
- Neutrino physics with Planck
  - Damping and phase shifts → Number of relativistic d.o.f
  - Gravitational lensing → Sum of neutrino masses

Removing lensing information send Planck parameters back to WMAP





Planck+WP LCDM best-fit

Planck+WP LCDM+Alens best-fit




Lensing info from power spectrum included
Planck+WP+highL
Planck+WP+highL (A_L)
Lensing info removed
P/P_{max}
1.0
0.8
0.6
0.4
0.2
0.0
0.0
0.4
0.8
1.2
1.6
2.0
Σm_ν[eV]


Lensing info from power spectrum included
Planck+WP+highL
Planck+WP+highL (A_L)
Planck+lensing+WP+highL
Lensing info removed
Lensing info from power spectrum and lensing reconstruction included
P/P_{max}
Σm_ν[eV]
0.0
0.2
0.4
0.6
0.8
1.0
0.0
0.4
0.8
1.2
1.6
2.0

Massive neutrino impact on BAO-H_0-CMB




Planck TT+lowP
Planck TT,TE,EE+lowP
+BAO
P/P_{max}
c^2_{eff}
0.0
0.2
0.4
0.6
0.8
1.0
0.27
0.30
0.33
0.36


Planck TT+lowP
Planck TT,TE,EE+lowP
+BAO
P/P_{max}
c^2_{vis}
0.0
0.2
0.4
0.6
0.8
1.0
0.00
0.25
0.50
0.75
1.00

Galactic Cleaning Cross-Check

$$T^{\mathrm{clean}} = (T^{\mathrm{freq}} - x\,T^{545})/(1-x)$$

- For the power spectrum analysis we use frequency maps which we clean in two ways
  - Model the power spectrum of the galaxy and subtract it (2013, 2014 baseline)
  - Subtract a high frequency map (“Mspec", new in 2014 for cross check since we use more sky)



Top row: Difference between
TT spectrum and CMB best-fit

Mspec (map cleaning)
Plik (2014 baseline)



Bottom row: after subtracting the foreground spectrum as well

Excellent power spectrum (and also cosmological parameter) agreement.
=> Galactic dust cleaning is under very good control.

TE & EE systematics

- The polarization power spectra are not yet sufficiently well tested to use in the main analysis.
- There are low-level systematics largely from to T->P leakage due to beam mismatch.
- This will be fixed for 2015



TE spectrum – best-fit CMB (from TT)

Outline

- Planck
- **ΛCDM, the standard model of cosmology, passes another precision test**
- Hands on with the Planck data
- Number of neutrino-like species $N_{\rm eff}$
  - BAO and H_0
  - Damping and acoustic oscillation phase shifts
  - Constraints on axions and axion-like particles (ALPs)
- Sum of neutrino masses Σm_ν
  - Gravitational lensing


(highly uncertain)
Key Epochs in the Standard Cosmological Model
Radiation
Matter
Dark energy
Inflation ends
opaque
nucleosynthesis
10-27 10-24 10-21 10-18 10-15 10-12 10-9 10-6 10-3 1
Scale Factor (a)


(highly uncertain)
Inflation ends
10^{-51} 10^{-48} 10^{-45} 10^{-42} 10^{-39} 10^{-36} 10^{-33} 10^{-30} 10^{-27} 10^{-24}
Scale Factor (a)


(highly uncertain)
Inflation ends
Density fluctuations created that
lead to observed CMB anisotropy.
10^{-51} 10^{-48} 10^{-45} 10^{-42} 10^{-39} 10^{-36} 10^{-33} 10^{-30} 10^{-27} 10^{-24}
Scale Factor (a)


(highly uncertain)
Inflation ends
A_s, n_s
Density fluctuations created that lead to observed CMB anisotropy.
10^{-51} 10^{-48} 10^{-45} 10^{-42} 10^{-39} 10^{-36} 10^{-33} 10^{-30} 10^{-27} 10^{-24}
Scale Factor (a)


(highly uncertain)
Inflation ends
Radiation
Matter
Dark energy
opaque
nucleosynthesis
10-27
10-24
10-21
10-18
10-15
10-12
10-9
10-6
10-3
1
Scale Factor (a)


(highly uncertain)
Inflation ends
Radiation
Matter
ρ_γ, ρ_ν
Dark energy
opaque
nucleosynthesis
10^{-27} 10^{-24} 10^{-21} 10^{-18} 10^{-15} 10^{-12} 10^{-9} 10^{-6} 10^{-3} 1
Scale Factor (a)


(highly uncertain)
Inflation ends
Radiation
Matter
ρ_c, ρ_b
ρ_Λ
Dark energy
opaque
$a_{\rm reion}$
nucleosynthesis
10^{-27} 10^{-24} 10^{-21} 10^{-18} 10^{-15} 10^{-12} 10^{-9} 10^{-6} 10^{-3} 1
Scale Factor (a)

The six-parameter ΛCDM model

A_s, n_s

Governs Spectrum of Primordial fluctuations

$$\rho_b = \Omega_b h^2$$
$$\rho_c = \Omega_c h^2$$
$$\rho_\Lambda = \Omega_\Lambda h^2$$

Matter Content

Optical depth to reionization

$a_{\rm reion} \rightarrow \tau$

The six-parameter ΛCDM model

A_s, n_s

Governs Spectrum of Primordial fluctuations

$$\rho_b = \Omega_b h^2$$
$$\rho_c = \Omega_c h^2$$
$$\rho_\Lambda = \Omega_\Lambda h^2$$

Matter Content

$N_{\rm eff}, \Sigma m_\nu, w, \Omega_K, ...$

Extensions

Optical depth to reionization

$a_{\rm reion} \rightarrow \tau$


0.35
0.30
0.25
0.20
0.15
Y_p
1
2
3
4
5
6
$N_{\rm eff}^{\rm CMB}$
6.75
6.60
6.45
6.30
6.15
6.00
5.85
5.70
5.55
$\log_{10} m_\phi / {\rm eV}$

The orbit

Planck makes a map of the full sky every ~6 months.